

PEEL
HERE

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



ACTIVE CONSTITUENT: 500g/L CHLOROTHALONIL

GROUP **M5** FUNGICIDE

For the control of fungal diseases on almonds, apricots, bananas, carrots, celery, cherries, faba beans, grapes, onions, peaches, peanuts, peas, plums, potatoes, tomatoes and vegetables as per the Directions for Use Table.

IMPORTANT: READ THIS LEAFLET BEFORE USING THIS PRODUCT

PROTECTION OF WILDLIFE, FISH, CUSTACEA AND ENVIRONMENT

Dangerous to fish. Do NOT contaminate streams, rivers or waterways, with chemical or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If not recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

SAFETY DIRECTIONS

Attacks eyes and skin. Avoid contact with eyes and skin. Do not inhale spray mist. Repeated exposure may cause allergic disorders. Sensitive workers should wear protective clothing. When preparing spray and using prepared spray wear cotton overalls buttoned to the neck and wrist, washable hat, elbow-length PVC gloves and goggles and disposable fume mask. If clothing becomes contaminated with product or wet with spray remove clothing immediately. If product on skin immediately wash area with soap and water. If product in eyes, wash out immediately with water. After use and before eating, drinking or smoking wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, goggles and contaminated clothing.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre. Telephone: Australia 13 11 26. If swallowed, do NOT induce vomiting. Give a glass of water.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet which can be obtained from the supplier

LIMIT OF WARRANTY AND LIABILITY

Axichem Pty Ltd warrants that this material conforms to the chemical description on the label. As the use of product sold is beyond the control of Axichem Pty Ltd, no responsibility whatsoever for any consequences is accepted in respect of this product, save those non-excludable conditions implied by any State and Federal legislation or law of a Territory. Not for repackaging or reformulations. No licence under any non-Australian patent is granted or implied by purchase of this container.

APVMA Approval No: 81995/104774

Batch Number:

Date of Manufacture:

IN A TRANSPORT EMERGENCY DIAL 000 POLICE OR FIRE BRIGADE	FOR SPECIALIST ADVICE IN AN EMERGENCY ONLY CALL 1800 039 008 ALL HOURS AUSTRALIA WIDE
--	---

AxiChem
Pty Ltd

AXICHEM PTY LTD
A.B.N 35 131 628 594
9 PALINGS COURT, NERANG QLD 4211
PHONE: 07 5596 1736 www.axichem.com.au

AxiChem

DIRECTIONS FOR USE**RESTRAINTS: DO NOT** mix spraying oils or wetting agents with this product.

Situation and Crop	Disease	Rate	WHP (Days)	Critical Comments	
Almonds (Not Qld, WA)	Shot hole (<i>Stigmia carpophila</i>) Stone fruit rust (<i>Tranzschella discolor</i>)	230mL/ 100L Apply 2000L/ha	-	Apply at bud-swell, bud burst, shuck fall, cap fall. Apply every 10-14 days. Apply 1 week pre-harvest.	
Apricots (Not Qld)	Brown rot- Fruit (<i>Monilinia fructicola</i>) Blossom blight (<i>Monilinia laxa</i>)		7	Apply at bud-swell, bud-burst, pink-bud, full bloom.	Apply no later than 35 days pre-harvest. See also additional restraints for stone fruits.
	Stone Fruit rust (<i>Tranzschella discolor</i>) Shot-hole (<i>Stigmia carpophila</i>), Freckle (<i>Venturia carpophila</i>)			Apply at bud-swell, bud burst, pink bud, shuck fall, cap fall, and then every 10-14 days.	
	Transit Rot (<i>Rhizopus stolonifer</i>)			Apply 7 days pre-harvest	
Bananas Do not apply to bananas within 14 days of applying an oil based spray or crop damage will occur.	Leaf Spot (<i>Mycosphaerella musicola</i>)	2.6L/ha (Qld, NSW, NT only)	1	For North Qld: Spray every 14 days late January to end of March. Extend to 21 days after March and then 28-35 days in the drier months of the year. For South Qld: Spray every 21 days from start of December to end of April. For NSW: Spray every 21 days from end of November to end of May. For WA: Spray every 10-14 days from November to end of May and extend to 14-21 days from May to November.	
		2.2L/ha (WA only)			
	Leaf speckle (<i>Mycosphaerella musae</i>)	4.3L/ha (Qld, NSW, WA, NT only)			Spray schedule same as for control of Leaf Spot. This product will not eradicate infection already present but will prevent the spread of disease.

Situation and Crop	Disease	Rate	WHP (Days)	Critical Comments
Broad Beans, Faba Beans (Tas only)	Chocolate spot (<i>Botrytis fabae</i>), Rust (<i>Uromyces viciae-fabae</i>)	1.6-2.6L/ha	7	Apply at the first sign of disease on the leaves or if weather conditions indicate likely incidence of disease. Repeat at 7-14 day intervals. Use the higher rate if the crop is dense and disease pressure is high.
Broccoli, Brussels Sprouts, Cabbages, Cauliflowers	Ring spot (<i>Mycosphaerella brassicicola</i>)	2.5-5L/ha		Apply at the first sign of disease on the leaves or if weather conditions indicate likely incidence of disease. Repeat at 14 day intervals. Use the higher rate if the crop is dense and disease pressure is high.
Carrots	Leaf Blight (<i>Alternaria dauci</i>)	2.6L/ha Or 230mL/100L		Commence application at first sign of disease and then spray every 7-14 days.
Cucurbits including Cantaloupe, Cucumbers, Honeydew, Muskmelon, Pumpkin, Squash, Rockmelon, Watermelon, Zucchini	Downy mildew (<i>Pseudoperonospora cubensis</i>), Gummy Stem Blight (<i>Didymella bryoniae</i>), Anthracnose (<i>Colletotrichum orbiculare</i>), Leaf Blight (<i>Alternaria cu</i>), Target Leaf Spot (<i>Cercospora citrullina</i>)	2.4-3.6L/ha	1	Apply on a 7-14 day schedule. Apply rates towards the higher end of the recommended range at 7-day intervals when disease pressure is high. Commence spray program early at first true leaf, and before vining or flowering. When used at these rates this product will suppress Powdery Mildew.
	Belly Rot (<i>Rhizoctonia solani</i>)			Suppression of belly rot will be obtained where fruit form on soil when sprayed using the above foliar spray programme.
Celery	Septoria leaf spot (<i>Septoria apicolica</i>)	2.6L/ha or 230mL/100L		Apply every 7-14 days. Apply also to seedbeds.
	Cercospora Early Blight (<i>Cercospora apii</i>) (NSW, ACT only)	1.8L/ha		Apply every 10-14 days.

Situation and Crop	Disease	Rate	WHP (Days)	Critical Comments
Cherries	Brown rot - Fruit (<i>Monilinia fructicola</i>) Blossom blight (NOT WA) (<i>Monilinia laxa</i>)	230mL/100L Apply 2000L/ha	7	Apply at bud-swell, bud-burst, pink bud, early blossom, full bloom. Apply 3 weeks pre-harvest and 1 week pre-harvest.
	Shot-hole <i>Stigmata carpophila</i> Stone fruit rust (<i>Tranzschelia discolor</i>)			Apply at bud-swell, bud burst, pink bud, shuck fall, cap fall and then apply every 10-14 days. Apply 1 week pre-harvest.
	Transit Rot (<i>Rhizopus stolonifer</i>)			Apply 1 week pre-harvest.
Grapes: Note: russetting of some table grape varieties may occur (Not WA)	Downy mildew (<i>Plasmopara viticola</i>) Bunch rot (<i>Botrytis cinerea</i>)	2.6-3.3L/ 100L or 230-300mL/100L	Dessert 7 Wine 14	Spray at the first appearance of the foliage disease. Apply at 7-14 day intervals particularly during wet weather and periods of rapid growth. For Bunch Rot, applications at early full bloom, two weeks later, at veraisson and two weeks prior to harvest. Table grapes- russetting of some varieties may occur. DO NOT apply after cap fall on these varieties.
Nectarines	Shot-hole (<i>Stigmata carpophila</i>) (NOT Qld, NSW)	230mL/100L Apply 2000L/ha	7	Apply at bud-swell, pink bud, shuck fall and then repeat each 14 days if weather conditions favour disease. Apply no later than 35 days pre-harvest. See also Additional Restraints for stone fruits.
	Brown rot -Fruit (<i>Monilinia fructicola</i>) Blossom blight (<i>Monilinia laxa</i>) (Tas only)			Apply at first bud-swell, bud burst, pink bud, early blossom and full bloom. Apply no later than 35 days pre-harvest. See also Additional Restraints for stone fruits.
Onions (excluding spring onions)	Downy Mildew (<i>Peronospora destructor</i>)	3.3L/ha		Apply at the first sign of disease on the leaves or if weather conditions indicate the likely incidence of disease. Repeat at 14 day intervals.

Situation and Crop	Disease	Rate	WHP (Days)	Critical Comments	
Ornamentals Chrysanthemums, Geraniums, Iris and other ornamentals	Botrytis, Grey mould, Rust diseases, <i>Septoria</i> and <i>Alternaria</i> leaf spot diseases (Tas only)	150mL/100L	-	Apply at 7-14 day intervals. Use 7 day interval under severe conditions. Avoid application to blooms in Azalea, Hydrangea and Petunia.	
Peaches	Brown rot -Fruit (<i>Monilinia fructicola</i>) Blossom blight (<i>Monilinia laxa</i>) (NOT Tas, WA)	230mL/100L Apply 2000L/ha	7	Apply at bud-swell, pink bud, early blossom and full bloom.	Apply no later than 35 days pre-harvest. See also Additional Restrains for stone fruits.
	Shot-hole (<i>Stigmia carpophila</i>) Stone fruit rust (<i>Tranzschelia discolor</i>)			Apply at bud-swell and continue at 7-14 day intervals.	
	Leaf curl (<i>Taphrina deformans</i>)			Apply at bud-swell and continue at 7-14 day intervals. In Qld spray only at bud-swell.	
Peanuts (NOT Vic, TAS, SA)	Early leaf spot (<i>Cercospora arachidicola</i>), Late leaf spot (<i>Cerosporidium personalum</i>), Peppery leaf spot (<i>Leplosphaerulina trifolii</i>), Peanut Rust (<i>Puccinia arachidis</i>), Net blotch (<i>Didymosphaeria arachildicola</i>)	1.6-2.6L/ha or 140-230mL/100L	-	Use higher rate for when conditions favour leaf spot or when rust appears. DO NOT feed peanut vines to livestock. NSW and WA only: Commence application when disease first appears and repeat as necessary to aintain control. QLD ONLY: North and South Burnett – Commence spraying when disease appears and repeat when rain and warm moist nights are expected. Atherton Tableland: Commence spraying four to six weeks after planting and repeat at 14-day intervals depending on conditions.	

Situation and Crop	Disease	Rate	WHP (Days)	Critical Comments	
Peas (Tas only)	Downey mildew (<i>Peronospora viciae</i>)	140-230mL/100L or 1.6-2.6L/ha	7	Apply at first sign of disease on the leaves or if weather conditions indicate likely incidence of disease. Repeat at 7-14 days intervals. Use the higher rate if the crop is dense and disease pressure is high.	
Plums	Brown rot -Fruit (<i>Monilinia fructicola</i>) Blossom blight (<i>Monilinia laxa</i>) (NOT WA)	230mL/ 100L Apply 2000L/ha	1	Apply at bud-swell, bud burst, early blossom, pink bud and full bloom.	Apply no later than 35 days pre-harvest. See also Additional Restraints for stone fruits.
	Shot-hole (<i>Stigmia carpophila</i>) Stone fruit rust (<i>Tranzschelia discolor</i>)			Apply at bud-swell, bud burst, shuck fall and cap fall.	
Potato	Early Blight or Target Spot (<i>Alternaria solani</i>) Late Blight or Irish Blight (<i>Phytophthora infestans</i>)	1.6-2.6L/ha	-	Spray at first sign of foliage disease or at flowering time, whichever is earlier. Spray every 7-14 days until within 14 days of desiccation or harvest.	
Rhubarb (NSW, Act only)	Assochyta leaf spot (<i>Ascochyta rhei</i>)	200mL/100L Apply 2000L/ha	1	Apply every 10-14 days.	
Tobacco (Qld only)	Tobacco leaf spot (<i>Ascochyta arida</i>)	320mL/100L	-	Apply every 7 days in seedbed application only.	
Tomato	Early Blight or Target Spot (<i>Alternaria solani</i>), Leaf Mould (<i>Fulvia fulva</i>), Grey Leaf Spot (<i>Stemphylium solari</i>) Botrytis Rot or Grey Mould (<i>Botrytis cinerea</i>)	2.6-3.3L/ha or 230-300mL/100L	1	Spray at first appearance of the foliage disease. Spray at 7-14 day intervals. Continue into the picking season to obtain best results against leaf mould and Botrytis rot.	
Turf (NSW only)	Grey leaf spot (<i>Pyricularia grisea</i>)	35L/ha	-	Apply to the leaf as a curative application. Do not irrigate for 24 hours after application.	

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED LEGISLATION.

WITHHOLDING PERIODS:

HARVEST

BANANAS, CELERY, CUCURBITS, PLUMS, RHUBARB, TOMATOES:

DO NOT HARVEST FOR 1 DAY AFTER APPLICATION.

BROCCOLI, BRUSSELS SPROUTS, CABBAGES, CAULIFLOWERS:

DO NOT HARVEST FOR 7 DAYS AFTER APPLICATION.

APRICOTS, CARROTS, CHERRIES, NECTARINES, PEACHES, TABLE GRAPES, FABA BEANS, ONIONS, PEAS:

DO NOT HARVEST FOR 7 DAYS AFTER APPLICATION.

WINE GRAPES: DO NOT HARVEST FOR 14 DAYS AFTER APPLICATION.

GRAZING

BEANS, PEANUTS, PEAS: DO NOT GRAZE LIVESTOCK ON TREATED CROPS.

POTATOES, PEANUTS (HARVEST): NOT REQUIRED WHEN USED AS DIRECTED

ADDITIONAL RESTRAINTS-STONE FRUITS

For all stone fruit except cherries, the following additional restraints apply:

Do not use oils after bud-swell and during the season AC Capsize 500 SC Fungicide is to be used after shuck fall.

Do not tank mix AC Capsize 500 SC Fungicide with EC formulations when spraying after shuck fall.

Do not exceed 4.6 L of AC Capsize 500 SC Fungicide per hectare of crop.

GENERAL INSTRUCTIONS

This product is a protectant fungicide for the control of a range of diseases in fruit and vegetables. For effective disease control a good leaf cover must be achieved before a widespread disease outbreak occurs.

Mixing:

Thoroughly shake and invert the pack several times before opening. Half fill spray vat and with agitation system running, add the required amount of product.

Application:

Low Volume - 200 or more litres per hectare.

High Volume - 500 - 1000 litres per hectare.

Aerial Application - 20 litres or more per hectare.

Mister - 75 - 255 litres per hectare.

The volume of water applied should increase as the plants grow in size to maintain adequate cover.

Compatibility:

This product is compatible with wettable powder and flowable formulations of most commonly used fungicides, insecticides and miticides. Unless experience in your area has shown other combinations to be physically compatible and non-injurious to your crop do not combine with other chemicals, especially emulsifiable concentrates.

Wetting Agents:

The addition of surfactants does not improve this product's efficiency and under some conditions may cause plant injury.

Fungicide Resistance Warning

GROUP	M5	FUNGICIDE
--------------	-----------	------------------

AC Capsize 500 SC Fungicide is a member of the multi-site activity group of fungicides. For fungicide resistance management the product is a Group M5 fungicide. Some naturally occurring individual fungi resistant to AC Capsize 500 SC Fungicide and other Group M5 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by AC Capsize 500 SC Fungicide or other Group M5 fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, Axichem Pty Limited accepts no liability for any losses that may result from the failure of AC Capsize 500 SC Fungicide to control resistant fungi.

PROTECTION OF WILDLIFE, FISH, CRUSTACEA AND ENVIRONMENT

Dangerous to fish. Do NOT contaminate streams, rivers or waterways with chemical or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

SAFETY DIRECTIONS

Attacks eyes and skin. Avoid contact with eyes and skin. Do not inhale spray mist. Repeated exposure may cause allergic disorders. Sensitive workers should wear protective clothing. When preparing spray and using prepared spray wear cotton overalls buttoned to the neck and wrist, washable hat, elbow-length PVC gloves and goggles and disposable fume mask. If clothing becomes contaminated with product or wet with spray remove clothing immediately. If product on skin immediately wash area with soap and water. If product in eyes, wash out immediately with water. After use and before eating, drinking or smoking wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, goggles and contaminated clothing.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre. Telephone: Australia 13 11 26. If swallowed, do NOT induce vomiting. Give a glass of water.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet which can be obtained from the supplier.

LABEL ELEMENTS

GHS Label Elements	
---------------------------	--

SIGNAL WORD	DANGER
--------------------	---------------

Hazard Statement(s)	
AUH066	Repeated exposure may cause skin dryness and cracking.
H317	May cause an allergic skin reaction.
H318	Causes serious eye damage.
H330	Fatal if inhaled.
H335	May cause respiratory irritation.
H351	Suspected of causing cancer.
H410	Very toxic to aquatic life with long lasting effects.

Precautionary statement(s) Prevention	
P201	Obtain special instructions before use.
P260	Do not breathe mist/vapours/spray.
P271	Use only outdoor or in a well-ventilated area.
P280	Wear protective gloves, protective clothing, eye protection and face protection.
P273	Avoid release to the environment.
P284	(In case of inadequate ventilation) wear respiratory protection.
P272	Contaminated work clothing should not be allowed out of the workplace.

Precautionary statement(s) Response	
P304+P340	IF INHALED: Remove person to fresh air and keep comfortable for breathing.
P305+P351+P338	IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.
P308+P313	IF exposed or concerned: Get medical advice/attention.
P310	Immediately call a POISON CENTRE/doctor/physician/first aider.
P302+P352	IF ON SKIN: Wash with plenty of water.
P333+P313	If skin irritation or rash occurs: Get medical advice/attention.

P362+P364	Take off contaminated clothing and wash it before reuse.
------------------	--

P391	Collect spillage.
-------------	-------------------

Precautionary statement(s) Storage

P403+P233	Store in a well-ventilated place. Keep container tightly closed.
------------------	--

P405	Store locked up.
-------------	------------------

Precautionary statement(s) Disposal

P501	Dispose of contents/container to authorised hazardous or special waste collection point in accordance with any local regulation.
-------------	--

LIMIT OF WARRANTY AND LIABILITY

Axichem Pty Ltd warrants that this material conforms to the chemical description on the label. As the use of product sold is beyond the control of Axichem Pty Ltd, no responsibility whatsoever for any consequences is accepted in respect of this product, save those non-excludable conditions implied by any State and Federal legislation or law of a Territory. Not for repacking or reformulations. No licence under any non-Australian patent is granted or implied by purchase of this container.

AxiChem

AxiChem